

UNIVERSITY OF CALIFORNIA, SANTA BARBARA
Department of Physics

Physics 110A
Prof. M.S. Witherell

December 9, 1991

FINAL

Melinda Kellogg

SHOW YOUR WORK AS CLEARLY AS POSSIBLE

The exam is closed book and notes, but you may use any information you have brought with you on one side of a $8\frac{1}{2}'' \times 11''$ sheet of paper. Do not spend too much time on one problem. Be sure to give answers in terms of given quantities.

#	pts
1	15
2	15
3	11
4	12
5	14
6	14
7	15
<hr/>	
	96

~~100~~

Good Work!

COURSE A+

1. A sphere of radius a contains charge of constant density:

$$\begin{aligned} \rho &= \rho_0 & r < a \\ \rho &= 0 & r > a \end{aligned}$$

a) Find ϕ everywhere.

15

b) Calculate the electrical energy stored, using $U_e = \frac{1}{2} \int \rho \phi d\tau$.

c) Calculate the electrical energy stored, using $U_e = \frac{1}{2} \int \epsilon_0 E^2 d\tau$.



a) $r > a$

$$\phi = \frac{Q_{in}}{4\pi\epsilon_0 r} = \frac{\rho_0 a^3}{3\epsilon_0 r}$$

5 $Q_{in} = \frac{4}{3}\pi a^3 \rho_0$

$r < a$

$$\vec{E} = \frac{Q_{in}}{4\pi\epsilon_0 r^2}$$

$$\vec{E} = \frac{\rho_0 r}{3\epsilon_0} \quad \phi(a) = \frac{\rho_0 a^2}{3\epsilon_0}$$

$$\phi(r) = \int_r^a \vec{E} \cdot d\vec{r} + \phi(a) = \frac{\rho_0}{3\epsilon_0} \int_r^a r dr = \frac{\rho_0}{6\epsilon_0} r^2 \Big|_r^a = \frac{\rho_0}{6\epsilon_0} (a^2 - r^2)$$

$$\Rightarrow \phi(r) = \frac{\rho_0}{6\epsilon_0} (a^2 - r^2) + \frac{\rho_0 a^2}{3\epsilon_0}$$

$$\frac{1}{3} - \frac{1}{5} = \frac{5}{15} - \frac{3}{15} = \frac{2}{15}$$

5 b) $U_e = \frac{1}{2} \int \rho \phi d\tau = \frac{1}{2} \rho_0 \int_0^a \int_0^{2\pi} \int_0^\pi \left(\frac{\rho_0}{6\epsilon_0} (a^2 - r^2) + \frac{\rho_0 a^2}{3\epsilon_0} \right) r^2 dr d\cos\theta d\phi$

$$= \frac{4\pi\rho_0^2}{2 \cdot 6\epsilon_0} \int_0^a (a^2 - r^2) r^2 dr + \frac{4\pi\rho_0^2 a^2}{6\epsilon_0} \int_0^a r^2 dr = \frac{4\pi\rho_0^2}{12\epsilon_0} \left[\frac{a^2 r^3}{3} \Big|_0^a - \frac{r^5}{5} \Big|_0^a \right] + \frac{4\pi\rho_0^2 a^2}{6\epsilon_0} \left[\frac{r^3}{3} \Big|_0^a \right]$$

$$= \frac{4\pi\rho_0^2}{12\epsilon_0} \left[\frac{a^5}{3} - \frac{a^5}{5} \right] + \frac{4\pi\rho_0^2 a^5}{6\epsilon_0 \cdot 3} = \frac{\rho_0^2 a^5 \pi}{\epsilon_0} \left(\frac{8}{12 \cdot 15} + \frac{4}{18 \cdot 9 \cdot 5} \right) = \frac{12}{45} \frac{\rho_0^2 a^5 \pi}{\epsilon_0}$$

$$\frac{12}{45}$$

Work Space

$$5) c) U_e = \frac{1}{2} \epsilon_0 \int_V E^2 d\tau$$

$$r > a \quad \vec{E} = \frac{\rho_0 a^3}{3\epsilon_0 r^2}$$


$$r < a \quad \vec{E} = \frac{\rho_0 r}{3\epsilon_0}$$

$$= \frac{1}{2} \epsilon_0 \int_0^{2\pi} \int_0^\pi \int_0^a \left(\frac{\rho_0 r}{3\epsilon_0}\right)^2 r^2 dr d\cos\theta d\phi + \frac{1}{2} \epsilon_0 \int_0^{2\pi} \int_0^\pi \int_a^\infty \left(\frac{\rho_0 a^3}{3\epsilon_0 r^2}\right)^2 r^2 dr d\cos\theta d\phi$$

$$= \frac{2\pi \rho_0^2}{9\epsilon_0} \int_0^a r^4 dr + \frac{2\pi \rho_0^2 a^6}{9\epsilon_0} \int_a^\infty \frac{1}{r^2} dr \quad r^2 \quad -\frac{1}{r} \Big|_a^\infty = \frac{1}{a}$$

$$= \frac{2\pi \rho_0^2 a^5}{45\epsilon_0} + \frac{5 \cdot 2\pi \rho_0^2 a^5}{5 \cdot 9\epsilon_0} = \boxed{\frac{12}{45} \frac{\rho_0^2 a^5 \pi}{\epsilon_0}}$$

Kelly



2. A coaxial cable consists of two conductors, one with outer radius a and the second with inner radius $b > a$, outer radius c . The gap between the conductors ($a < \rho < b$) is filled with a dielectric with dielectric constant κ_e . The inner conductor carries linear charge density λ , the outer carries density $-\lambda$.

- a) What is \vec{E} between the conductors?
- b) What is the capacitance per unit length?
- c) What is the volume bound charge density in the dielectric (ρ_b)?
- d) What is the surface bound charge density, σ_b , on the outer surface of the dielectric?
- e) What is the energy stored per unit length?

15

3 a) $\vec{D} = \frac{\lambda}{2\pi\rho} \hat{\rho} \Rightarrow \vec{E} = \frac{\vec{D}}{\kappa_e \epsilon_0} = \boxed{\frac{\lambda}{2\pi\epsilon_0 \rho \kappa_e} \hat{\rho}}$ ✓

3 b) $C = \frac{Q}{\Delta\phi}$ $\Delta\phi = + \int_a^b E_{\rho} d\rho = + \frac{\lambda}{2\pi\epsilon_0 \kappa_e} \int_a^b \frac{d\rho}{\rho} = + \frac{\lambda}{2\pi\epsilon_0 \kappa_e} \ln \frac{b}{a}$
 $Q = \lambda L$

$C = \frac{\lambda L 2\pi\epsilon_0 \kappa_e}{\lambda \ln(b/a)} = \boxed{\frac{2\pi\epsilon_0 \kappa_e}{\ln(b/a)}}$ ✓

3 c) $\vec{P} = (\kappa_e - 1)\epsilon_0 \vec{E} = \frac{(\kappa_e - 1)\lambda}{\kappa_e} \frac{1}{2\pi\rho} \hat{\rho}$

$\rho_b = -\vec{\nabla} \cdot \vec{P} = -\frac{1}{\rho} \frac{\partial}{\partial \rho} (\rho P_{\rho}) = \boxed{0}$ ✓

3 d) $\sigma_b = \vec{P} \cdot \hat{n} = \boxed{\frac{(\kappa_e - 1)\lambda}{\kappa_e} \frac{1}{2\pi b}}$ ✓ inner sum 0?

3 e) $U_e = \frac{1}{2} \int_V \vec{D} \cdot \vec{E} d\vec{r}$ $\vec{D} = \frac{\lambda}{2\pi\rho} \hat{\rho}$ $\vec{E} = \frac{\lambda}{2\pi\epsilon_0 \rho \kappa_e} \hat{\rho}$
 $= \frac{\lambda^2}{8\pi^2 \epsilon_0 \kappa_e} \int_0^{2\pi} \int_0^L \int_a^b \frac{1}{\rho^2} \rho d\rho d\phi dz = \frac{L\lambda^2}{4\pi\epsilon_0 \kappa_e} \ln \rho \Big|_a^b = \boxed{\frac{\lambda^2 \ln(b/a)}{4\pi\epsilon_0 \kappa_e}}$ ✓

